

Hidden Topological Order in ^{23}Na ($F = 1$) Bose-Einstein Condensates

Fei Zhou

ITP, Utrecht University, Minnaert Building, Leuvenlaan 4, 3584 CE Utrecht, The Netherlands
(February 1, 2008)

We show the existence of a new hidden topological order in ^{23}Na ($F=1$) Bose-Einstein condensates (BEC) with antiferromagnetic interactions. Occurrence of this order is due to the confinement of hedgehogs in spin ordered BEC where a spin Josephson effect takes place. However, a topological long range order is also argued to coexist with a short range spin correlation, as a result of topological order from disorder.

^{23}Na ($F = 1$) BEC with antiferromagnetic interactions turn out to have many extremely fascinating properties [1–9]. One of them is the topology of the order parameter space. It was pointed out in previous works that ^{23}Na BEC in optical traps are in *Quantum Spin Nematic States*, with the order parameter space being $[S^1 \times S^2]/Z_2$ [8,9]. In this article we will report a new hidden long range order of pure topological nature and explore the possibility of having spin correlated states based on a topological consideration. To study spin correlated BEC of spin-1 interacting atoms, it is important to properly choose a set of collective coordinates. These variables are to completely characterize the nonlinear spin dynamics in the whole parameter space and to yield a simplest representation for two-body spin dependent scatterings. For this reason we employ a vector $\mathbf{n}$ living on a unit sphere for the description of the nonlinear spin dynamics in the presence of antiferromagnetic interactions. The low energy spin dynamics in the system can be mapped into an $o(3)$ nonlinear sigma model (NL σ M) [8,9],

$$\mathcal{L} = \frac{1}{2f}(\partial_\mu \mathbf{n})^2, \mathbf{n}^2 = 1, \\ f = (16\pi)^{1/2}(\rho\Delta a^3)^{1/6}, \Delta a = \frac{a_2 - a_0}{3} \quad (1)$$

$a_{0,2}$ are the scattering lengths in $F = 0, 2$ channels and ρ is the density. We also have introduced dimensionless length and time: $\mathbf{r} \rightarrow \mathbf{r}/\rho^{1/3}$, $it \rightarrow \tau/v_s\rho^{1/3}$, and $v_s = \sqrt{16\pi\hbar^2\Delta a\rho/M^2}$. Derivatives ∂_μ are defined as $(\partial_\tau, \partial_x, \partial_y, \partial_z)$. A reduction from $[S^1 \times S^2]/Z_2$ to S^2 is possible when Z_2 fields discussed in [9] are effectively frozen, or Z_2 strings are gapped. Moreover, the local coupling between the $U(1)$ phase and spin order involves higher derivative terms and becomes important only at a rather high frequency $\hbar\rho^{2/3}/2M$.

Following Eq.1, the energy of the system consists of two parts: a) the potential energy $\hbar^2\rho(\nabla\mathbf{n})^2/2M$, which is the energy cost in the presence of a slow variation of $\mathbf{n}$, b) and the zero point kinetic (rotation) energy $I_0(\partial_t\mathbf{n})^2/2$, where $I_0 = (\hbar^2\Delta a\rho/M)^{-1}$ can be considered as the effective inertial of an individual atom. This inertial originates from two-body scatterings and is inversely proportional to scattering lengths. Zero point rotations tend to disrupt the order of $\mathbf{n}$ between different atoms. f is

a measure of the amplitude of quantum fluctuations in ^{23}Na BEC.

In weakly interacting limit f is much less than unity, the symmetry is broken in the ground state and the BEC have a spin order, with $\mathbf{n}(\mathbf{r}, t) = \mathbf{n}_0$. Just as the Josephson effect occurs in a superfluid where a phase long range order is established, a spin Josephson effect occurs in spin ordered BEC. A spinor condensate in a weak external magnetic field $B\mathbf{e}_z$ exhibits the following time dependence:

$$\Psi(t) = \frac{1}{\sqrt{2}}[\exp(i\mu_+t)|1, 1\rangle + \exp(i\mu_-t)|1, -1\rangle], \quad (2)$$

and $\mu_\pm = \mu_1 \pm \omega_B$, $\omega_B = g\mu_B B$ (g is the Lande factor of atoms). Notice that in the presence of a spin stiffness, a well defined relative phase between two components of the condensates is developed. To observe the spin coherence, one introduces a reference condensate which is at a chemical potential μ_2 but experiences no magnetic fields. The temporal dependence of the interference between these two condensates would be

$$\delta\rho(t) \propto \cos((\mu_1 - \mu_2)t) \cos(\omega_B t). \quad (3)$$

At $\omega_B = 0$, Eq. 3 reflects to a usual AC Josephson effect and when $\mu_1 = \mu_2$, one observes an AC spin Josephson effect due to a spin order in BEC, a sinusoidal-time dependence with frequency determined by ω_B instead of difference in chemical potentials. The alternative way to observe the spin coherence without introducing a reference condensate is to transform $|1, 1\rangle$ component into $|1, -1\rangle$ after the magnetic field is switched on for time t . This results in an interference between the induced $|1, -1\rangle$ component and the original $|1, -1\rangle$ component. The condensate density in the overlapped region exhibits a sinusoidal dependence on the waiting time t with a frequency $2\omega_B$.

Quantum fluctuations of the nematic order can be estimated in a lowest order approximation. Introducing $\delta\mathbf{n} = \mathbf{n} - \mathbf{n}_0$ and assuming the fluctuations are weak, we obtain, $\langle (\delta\mathbf{n}^2) \rangle \propto f$. As f increases, spin wave excitations start to interact strongly and Eq.2 becomes invalid. The renormalization group (RG) equation of f is determined by the interactions between collective modes. In $3 + 1$ dimension, the RG equation

takes a form $df/dl = \beta(f)$, with the β -function given as $\beta(f) = -2f(f_c - f)$. Within the frame work of NL σ M, $f_c = 8\pi^2$ in $d = 3$. f characterizes spin correlations in the ground state of ^{23}Na BEC.

At a high density limit $f > f_c$ [10], zero point kinetic energy dominates and the spin stiffness is renormalized to zero at a long wave length limit. Especially, in an extremely quantum disordered phase, the potential energy at interatomic scale $E_T = \hbar^2 \rho^{2/3}/2M$ is negligible compared with zero point rotation energy $E_o = \hbar^2/2I_0$ of an individual atom. $\mathbf{n}$ of each atom fluctuates independently and spins of atoms only correlate at an interatomic distance. The excitation spectrum has a gap of order E_o . As the two-body scattering gets weaker, the inertial of each individual atom I_0 is higher and the motion of $\mathbf{n}$ on the unit sphere becomes slower. Thus, $\mathbf{n}$ of different atoms starts to correlate at a finite length ξ , which is a function of the parameter f , much longer than the interatomic distance. Under the current situation, $\xi = \rho^{-1/3} f_c/[f - f_c]$. And the state is rotation invariant. The temporal correlation is characterized by τ_ξ , the time needed for a spin wave to propagate over a distance ξ with a velocity v_s . At a low density limit $f < f_c$, the potential energy dominates and the spin stiffness flows to a finite value under the renormalization group transformation. All spin-1 Bosons rotate as a rigid body with an effective inertial $I(N) = NI_0$. The energy gap in the excitation spectrum vanishes $E_{gap} \propto \hbar^2/2NI_0$ as the number of Bosons N approaches infinity, signifying the existence of gapless Goldstone mode.

For ^{23}Na BEC confined in highly anisotropic traps, zero point motions of collective variables are more prominent. Let us approximate highly anisotropic traps as 1D systems. The β -function RG equation in this case can be shown as $\beta(f) = 1/2\pi f^2$, which always flows into a strong coupling point, i.e. a rotation invariant state. And the correlation length is given as $\xi = \rho^{-1} \exp(4\pi/f)$ (ρ is the linear density). However, the length of traps has to be longer than ξ in order for this spin liquid to be observed in experiments.

To study the topological long range order, we introduce a connection field to characterize all configurations of $\mathbf{n}$ [9]. The field strength is a gauge invariant Pontryagin density $\mathbf{F}_{\mu\nu} = \epsilon^{\mu\nu} \epsilon^{\alpha\beta\gamma} \mathbf{n}_\alpha \partial_\mu \mathbf{n}_\beta \partial_\nu \mathbf{n}_\gamma$. In $(3+1)D$, $\mathbf{F}_{0i}$, $i = x, y, z$ corresponds to an electric field and $\epsilon^{ijk} \mathbf{F}_{ij} = \mathbf{H}_k$ represents three components of magnetic field; in $2d$, $\mathbf{F}_{0i}$, $i = x, y$ is an electric field with two components and the magnetic field has only z component $\mathbf{H}_z = \mathbf{F}_{xy}$. Particularly, we will be interested in the fluctuations of a topological charge defined as

$$C_m = \frac{1}{4\pi} \oint_s d\mathbf{S} \mathbf{H} \cdot \mathbf{e}_n, \quad (4)$$

where the integral is carried over the boundary of volume $\mathcal{V}$ and $\mathbf{e}_n$ is a unit vector normal to the boundary. We are going to show that at large $\mathcal{V}, t$ limit,

$$\lim_{t, \mathcal{V} \rightarrow \infty} \langle (C_m(t) - C_m(0))^2 \rangle = 0 \quad (5)$$

in both spin ordered and spin weakly disordered BEC, as results of a topological long range order. However, in strongly disordered limit,

$$\lim_{t, \mathcal{V} \rightarrow \infty} \langle (C_m(t) - C_m(0))^2 \rangle = \frac{t\mathcal{V}}{\tau_m} \quad (6)$$

where τ_m is finite depending on tunneling rate of magnetic monopoles (see below). Averages in Eqs.5,6 are taken over the many-spin wavefunction $\Psi(\{\mathbf{n}\})$. In 2D, we define $C_m = (4\pi)^{-1} \int dx dy \mathbf{H}_z$, the total number of Skyrmions living on the 2D sheet and Eqs.5,6 hold in spin ordered and spin disordered BEC respectively ($\mathcal{V}$ is replaced by an area $\mathcal{S}$). The topological long range order occurs when hedgehogs are confined, either due to the condensation of atoms or due to an order from disorder mechanism.

Let us start with 2D. First, we notice that in space $\mathbf{x} = (\tau, \mathbf{r})$, $\mathbf{H}_\eta = \epsilon^{\eta\mu\nu} \mathbf{F}_{\mu\nu}$ is the solution of the Gauss equation

$$\partial_\eta \mathbf{H}_\eta(\mathbf{x}) = \sum_m Q_m \delta(\mathbf{x} - \mathbf{x}^m) \quad (7)$$

in the presence of space-time monopoles $\{\mathbf{x}^m\}$ in $(2+1)d$ Euclidean space.




Fig.1. Spin configurations at a) $t = +\infty$ and b) $t = -\infty$ boundaries of a monopole in $(2+1)d$.

The space-time monopoles represent instantons, which essentially connect a trivial vacuum to a Skyrmion configuration, as seen in Fig.1. In Euclidean space, a space-time monopole located at $x = (\tau_0, \mathbf{r}_0)$ terminates a Skyrmion centered at $\mathbf{r}_0$ at time τ_0 . Each monopole causes a change in topological charge C_m by one unit; the topological density $C_m = 1$ is conserved in the absence of space-time monopoles in $(2+1)D$. Indeed, following Eqs.4,7,

$$\frac{\partial C_m(\tau)}{\partial \tau} = \sum Q_m \delta(\tau - \tau_m), \quad (8)$$

where the surface contribution has been neglected since we will be interested in C_m per unit volume.

The probability of finding deconfined space-time monopoles depends on the energy of Skyrmions with $C_m = 1$ with respect to $C_m = 0$ configuration. In polar coordinates (r, ϕ) , a static Skymion is a configuration with $\mathbf{n}(\rho, \phi) = (\sin \theta(r) \cos \phi, \sin \theta(r) \sin \phi, \cos \theta(r))$; $\theta(r)$ varies from 0 at $r = 0$ to π at $r \gg \xi$, with ξ an arbitrary parameter. For spin ordered BEC where spin Josephson effects should be observed, the energy of the Skymion is proportional to $8\pi\hbar^2\rho/2M$ and is scale invariant. For a Skymion of given ξ [12], the connection field is concentrated in a region of the size ξ , $\mathbf{H}_z = r^{-1} \sin \theta(r) \partial \theta(r) / \partial r$, and gets screened at a length scale larger than ξ . In spin ordered BEC, a Skymion is nondegenerate with respect to a trivial vacuum. As we will see, this leads to the confinement of space-time monopoles.

To illustrate the point of confinement, we consider C_m as a function of time at some discrete unit and find the following binary representation for monopoles.




Fig.2. In a binary representation, monopole-like instants are represented by kinks living on a string, with C_m , the topological charge as the order parameter.

Each bit shown in Fig.2 labeled as 0, or 1 represents the topological charge read out at certain moment; a 1-bit corresponds to a Skymion while a 0-bit is for a $C_m = 0$ configuration. The a) string represents a trivial vacuum where C_m remains zero at all time. In the second binary string b), a 1-string stands for any scale invariant Skymion created at the time t_0 defined by the domain wall separating 0-string and 1-string. So a domain wall in our binary representation represents a monopole $Q_m = 1$, terminating a Skymion configuration ($C_m = 1$) at the interface and changing C_m by one unit. A domain wall-anti domain wall pair in the c) line corresponds to a monopole-antimonopole pair with $Q_m = \pm 1$; a 1-string here, or a Skymion, is terminated at a monopole at one end t_1 and at an antimonopole at the other end t_2 .

Since the energy of a Skymion in spin ordered BEC is higher than that of a trivial configuration, the energy of a 1-bit is positive with respect to a 0-bit. The energy of a domain wall in Eq.1 which is the number of 1-bits in the structure, defines the action of the monopole and is proportional to the length of the 1-string. The action of an isolated space-time monopole, or creation of a Skymion therefore is proportional to $L_t \hbar^2 \rho / 2M$, being infinity. (L_t is the perimeter along a temporal direction.)

The action of having monopole-anti-monopole pairs is proportional to $(t_1 - t_2)$, the time interval between creation and annihilation of Skyrmions, thus monopoles are confined. For this reason, Eq.8 vanishes and a topological long range order prevails.

The conclusion arrived so far can be extended to 3D straightforwardly. The change of the topological density is due to the quantum nucleation of monopoles instead of Skyrmions. In the binary representation in Fig.2, a 1-bit stands for a static monopole and a 0-bit for a trivial configuration. A domain-wall represents a termination of a static monopole at certain time. Since the energy of a monopole is proportional to the system size L in spin ordered BEC, the monopole is nondegenerate with respect to the trivial vacuum. Following the same argument carried out in 2D, the action to have a monopole nucleated is proportional to $LL_t \hbar^2 \rho / 2M$, and is infinity. This result has a profound impact on the monopole confinement. The interaction between a monopole and an antimonopole has to be linear in terms of distance between them. The action of a monopole pair nucleated at t_1 and annihilated at t_2 with largest spatial separation $|r_1 - r_2|$ is proportional to $|r_1 - r_2|(t_1 - t_2)$ and monopoles are confined because of spin stiffness. Therefore the right hand side of Eq.8 has to be zero because of the confinement, and topological charge C_m is a conserved quantity. We once again arrive at Eq.5 in a spin ordered BEC and topological charge is conserved.

The situation in disordered limit is more delicate and depends on dimensionalities. In spin disordered BEC, the spin stiffness is renormalized to zero and spin fluctuations are gapped. In the absence of spin stiffness, the energy of a Skymion is determined by its interaction with spin fluctuations. Upon integration of spin wave excitations, based on a general consideration of the gauge invariance and the parity invariance, we conclude Eq.1 should be reduced to $\mathcal{L}_s(\mathbf{F}_{\mu\nu}) = \mathbf{F}_{\mu\nu} \mathbf{F}_{\mu\nu} / 2g + \dots$; $g^{-1}(\Delta_s)$ is a function of the spin gap Δ_s measured in units of $(E_T E_o)^{1/2} = \hbar^2 (\Delta a \rho^{5/3})^{1/2} / 2M$. In a leading approximation, $g^{-1}(x)$ is a logarithmic function of x , i.e. $\ln x^{-1}$ in 3D and is a linear function of x in 2D [13]. We will analysis the topological order based on $\mathcal{L}_s$. The energy of C_m -configuration is $\alpha(L)C_m^2$ if quantum tunneling is neglected, with $\alpha(L)$ a function of the system size.

In 2D, with induced interactions, Skymion energy is no longer scale invariant and is minimized at $\xi = \infty$. The connection field of a Skymion spreads over the whole 2D sheet. The energy of a Skymion scales as L^{-2} and vanishes as the system size L goes to infinity. This implies that skymion configurations become degenerate with a trivial vacuum and monopoles are deconfined. This mechanism of deconfinement resembles the liberation of fractionalized quasiparticles in one dimension polymers [11]. In one dimensional polyacetylene, the ground state

has twofold degeneracy because of the Peierls instability and domain wall solitons become free excitations.

When instantons are liberated, the energy of a configuration C_m is ill-defined. A more serious consideration beyond the mean field approach outlined above involves the evaluation of monopole-like instantons action. In $(2+1)d$ the partition function of a monopole configuration $\{\mathbf{x}^m\}$, is $\sum_n P^n/n! \exp(-S_{in})$. And $P = L^2 L_t \exp(-\pi/g)$ is the quantum tunneling amplitude; $S_{in} = g^{-1} \sum_{m' \neq m} \pi Q_m Q_{m'} / |\mathbf{x}^{m'} - \mathbf{x}^m|$ represents the Coulomb interaction between space-time monopoles. The above result suggests that Skyrmions always condense in the spin disordered BEC and topological charge C_m is not a conserved quantity. It is important to realize that in the absence of Z_2 strings, ± 1 Skyrmions can be physically and homotopically distinguished because of the coherence of *quantum* spin nematic BEC, unlike the situation in a *classical* nematic liquid crystal. By moving a Skyrmion around a π -disclination, the condensate acquires a π -phase with respect to the original one, which can manifest itself in a Josephson type of effect. One can also distinguish ± 1 Skyrmions by looking at the local connection field. In a positive Skyrmion configuration, a spin- $\frac{1}{2}$ collective excitation, which carries half charge with respect of connection fields, experiences a connection field of an opposite sign compared to that of a negative Skyrmion [14].

So we are able to show that the change of C_m has a short range temporal correlation because of instanton effects,

$$< \frac{\partial C_m(\tau)}{\partial \tau} \frac{\partial C_m(0)}{\partial \tau} > = \exp(-\frac{tS}{\tau_m}). \quad (9)$$

And τ_m is proportional to $\xi^2 \tau_\xi \exp(\pi/g)$. Immediately, one recognizes that Eq.9 simply indicates a random walk of C_m as a function of time in spin disordered 2D BEC. Therefore, we conclude Eq.6 holds.

But in 3D weakly disordered BEC where stiffness has vanished, a static monopole still carries a finite energy because of spin fluctuations induced interactions. In 3D, as illustrated in $\mathcal{L}_s(\mathbf{F}_{\mu\nu})$, spin fluctuations discriminate topologically different configurations. Particularly, the fluctuations are strongest in $C_m = 0$ configuration so that the energy is lowest in the topological trivial configuration. The situation differs from 2D spin disordered limit where the energy of topological nontrivial configuration with $C_m = 1$ is the same as that of $C_m = 0$ one. For a monopole, each spherical shell of radius R can be viewed to be a Skyrmion squeezed into a 2D sheet of size R . The smaller shells, or textures of a finite size, dominate the energy of a monopole. Spin fluctuations interacting with the singularity lift a degeneracy between the monopole configuration and a trivial vacuum in the absence of spin stiffness. It also appears to us that this is independent of the form of $\mathcal{L}_s(\mathbf{F}_{\mu\nu})$ we introduced. The

energy of a monopole here is inversely proportional to g , i.e. $(E_o E_T)^{1/2} \ln \Delta_s^{-1}$. The action of having a monopole, which leads to a change in C_m by one unit, is proportional to L_t and is infinite for this reason.

So the monopoles remain confined even after spin correlation becomes short ranged, with the action to have a pair of monopoles being proportional to $(t_1 - t_2)$. Only pair production which conserves C_m is allowed. The topological long range order thus coexists with a short range spin correlation in 3D. This can be considered as a case of order from disorder phenomena. And, the confinement of monopoles in this case is driven purely by the spin fluctuations, instead of spin stiffness in spin ordered BEC. However, by increasing the spin disorder further, the degeneracy between $C_m = 1$ and $C_m = 0$ could be established, which signifies deconfinement of monopoles and breakdown of the long range topological order.

It is also possible to demonstrate the topological order/disorder in terms of space-time correlations of connection fields or Wilson loop integrals. The topological order is vital for the quantum number fractionalization of excitations. In 3D spin disordered BEC, where a topological long range order is established, spin- $\frac{1}{2}$ excitations are elementary ones [14]. On the other hand, when hedgehogs are liberated or condense, only spin-1 excitations exist in the spectrum. Finally, quantum hidden orders in spin liquids of strongly correlated electrons are recently reviewed by Wen [15]. Topological orders in spin triplet superconducting liquids are also investigated by Demler et al. [16]. I would like to thank ASPEN center of Physics and Amsterdam summer workshop on "Flux, charge, topology and statistics" for their hospitalities. Discussions with A. Abanov, Duncan Haldane, N. Read, T. Senthil, O. Starykh and Paul Wiegmann are greatly acknowledged.

-
- [1] C. J. Myatt, et.al., Phys. Rev. Lett. **78**, 586(1997).
 - [2] D. M. Stamper-Kurn, et.al., Phys. Rev. Lett. **80**, 2027(1998).
 - [3] J. Stenger, et.al., Nature **396**, 345(1998).
 - [4] T. L. Ho, Phys. Rev. Lett. **81**, 742 (1998).
 - [5] T. Ohmi and K. Machinda, J. Phys. Soc. Jpn. **67**, 1822(1998).
 - [6] C. K. Law, et.al., Phys. Rev. Lett. **81**, 5257(1998).
 - [7] Y. Castin and C. Herzog, cond-mat/0012040.
 - [8] F. Zhou and F. D. Haldane, ITP-UU-00/51(2000); F. Zhou, cond-mat/0102237.
 - [9] E. Demler, F. Zhou and F. D. Haldane, ITP-UU-01/09(2001); E. Demler and F. Zhou, cond-mat/0104409.
 - [10] This limit is difficult to approach in single traps because of fast recombination processes between alkali atoms. However, the difficulty can be avoided in an optical lat-

tice where $f = \sqrt{E_o/E_T}$; E_o is the zero point rotation energy and E_T is the exchange energy. See discussions in Ref.9.

- [11] For an excellent review, see A. J. Heeger, S. Kivelson, W. P. Su and J. R. Schrieffer, Rev. Mod. Phys. **60**, 781(1988).
- [12] Additional high derivative terms usually stabilize Skyrmions at a finite ξ .
- [13] This ansatz can be justified in the context of CP^N fields,

in a large- N expansion. See A. M. Polyakov, *Gauge Fields and Strings*, Harwood academic publishers (1987).

- [14] Fei Zhou, cond-mat/0106133; ITP-UU-01/23(2001).
- [15] X. G. Wen, cond-mat/0107071. Examples of topological spin fluids in strongly correlated electron systems can also be found in P. Wiegmann, Phys. Rev. Lett. **B 60**, 821(1988); X. G. Wen, F. Wilczek and A. Zee, Phys. Rev. **B 39**, 11413(1990).
- [16] E. Demler et al., cond-mat/0105446.